



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Bo

Search PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Ann Thorac Surg. 1998 May;65(5):1313-5.

Related Articles

Comment in:

- Ann Thorac Surg. 1998 Dec;66(6):2158-9.

**ELSEVIER SCIENCE
FULL-TEXT ARTICLE****Aortic valve operations under deep hypothermic circulatory arr
for the porcelain aorta: "no-touch" technique.****Byrne JG, Aranki SF, Cohn LH.**Division of Cardiac Surgery, Brigham and Women's Hospital, Harvard Medi
School, Boston, Massachusetts 02115, USA.

BACKGROUND: Aortic valve replacement or repair becomes a high-risk procedure in patients in whom the ascending aorta cannot be clamped either because of extensive calcification and risk of cerebral embolus or because of extensive adhesions precluding safe dissection and clamping. **METHODS:** We report the results of aortic valve replacement or repair with deep hypothermic circulatory arrest in 3 patients. Techniques to improve results include routine of epiaortic and transesophageal echocardiography, avoidance of manipulation the ascending aorta until the circulation is arrested, avoidance of antegrade cardioplegia, routine use of retrograde cardioplegia and retrograde cerebral perfusion, when feasible, and minimal aortotomy (just enough to excise and replace or repair the valve). **RESULTS:** Operations were accomplished in approximately 1 hour each with minimal manipulation of the aorta, thus minimizing aortic trauma and subsequent risk of cerebral embolus. Each patient had an unremarkable recovery without neurologic complications. **CONCLUSIONS:** Aortic valve replacement or repair using the "no-touch" technique and deep hypothermic circulatory arrest is the preferred method when dealing with the porcelain or unclampable aorta.

PMID: 9594858 [PubMed - indexed for MEDLINE]

Display

Abstract

Show:

20

Sort

Send to

Text

[Write to the Help Desk](#)

NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

Jul 17 2003